

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

SEMIANNUAL PROGRESS REPORT

ON

NASA GRANT NAG-1-410

PROJECT TITLE: Construction of Finite Difference Schemes Having
Special Properties for Ordinary and Partial Differential Equations

(NASA-CR-173722) CONSTRUCTION OF FINITE
DIFFERENCE SCHEMES HAVING SPECIAL PROPERTIES
FOR ORDINARY AND PARTIAL DIFFERENTIAL
EQUATIONS Annual Progress Report (Atlanta
Univ., Ga.) 5 p HC A02/MF A01

N84-28544

Unclass
00987

CSCL 12A G3/64

PRINCIPAL INVESTIGATOR: Ronald E. Mickens
Professor of Physics
Atlanta University
Atlanta, Georgia 30314
(404) 681-0251, ext. 327 or 336

STARTING DATE OF GRANT: October 1, 1983

TECHNICAL MONITOR: Dr. John N. Shoosmith
NASA Langley Research Center
Mail Stop 125
Hampton, Virginia 23665
(804) 865-3466

DUE DATE: April 1, 1984

This report summarizes my recent work on the construction of finite-difference models of differential equations having zero truncation errors.¹ The details of the calculations which follow will not be given since a paper is being written on these topics.

I. Unidirectional Wave Equation

The unidirectional, linear wave equation in one-dimension is

$$(1) \quad u_x + u_t = \lambda u$$

$$\lambda = \text{constant.}$$

For the initial value problem, where

$$(2) \quad u(x,0) = f(x) = \text{given function,}$$

the exact, general solution is

$$(3) \quad u(x,t) = e^{\lambda t} f(x-t) \quad .$$

If we define

$$(4) \quad u(x_m, t_n) = u_m^n$$

$$x_m = (\Delta x)m \quad , \quad t_n = (\Delta t)n$$

where (m,n) are integers, then it is easy to prove that the following finite-difference scheme is a zero truncation error model of eq. (1):

$$(5a) \quad u_m^{n+1} = e^{\lambda h} u_{m-1}^n$$

$$\Delta x = \Delta t = h$$

where

$$(5b) \quad u_m^0 = f((\Delta x)m) \quad .$$

ORIGINAL PAGE IS
OF POOR QUALITY

It should be pointed out that the conventional use of either the Euler forward- or backward- difference schemes for either of the derivatives in eq. (1) does not lead to eq. (5a).² In more detail, the "full" difference scheme associated with eq. (1) which has zero truncation error is easily gotten from eq. (5a) and written out in detail is

$$(6) \quad \frac{\Delta_t^{(+)} u_m^n}{\left(\frac{e^{\lambda h} - 1}{\lambda} \right)} + \frac{\Delta_t^{(-)} u_m^n}{\left(\frac{e^{\lambda h} - 1}{\lambda} \right)} = \lambda u_{m-1}^n$$

where

$$(7a) \quad \Delta_t^{(+)} u_m^n = u_m^{n+1} - u_m^n$$

$$(7b) \quad \Delta_x^{(+)} u_m^n = u_m^n - u_{m-1}^n$$

Note that the conventional application of finite difference techniques to eq. (1) would give

$$(8) \quad \frac{\Delta_t^{(+)} u_m^n}{\Delta t} + \frac{\Delta_x^{(+)} u_m^n}{\Delta x} = \lambda u_m^n$$

or some similar result, which does have a non-zero truncation error. Direct comparison of eqs. (6) and (8) shows that previous knowledge of finite-difference models for differential equations may not be of great help in the construction of schemes with zero-truncation error.^{1,2}

For practical problems where eq. (1) is to be numerically integrated, either eq. (5a) or eq. (6) may be used since they are mathematical equivalent expressions. However, from a computational point of view, eq. (5a) should be used.

Recently, I have been able to obtain a zero truncation error difference model for the following nonlinear, uni-directional wave equation

$$(9) \quad u_t + u_x = \lambda u^n \quad (n = \text{positive integer}) \quad .$$

The general case is rather difficult to write down; however, for $n = 2$, we have

$$(10a) \quad u_t + u_x = \lambda u^2$$

and the difference scheme is

$$(10b) \quad u_m^{n+1} = \frac{u_{m-1}^n}{1 - (\lambda \Delta x) u_{m-1}^n} \quad .$$

where

$$(10c) \quad \Delta t = \Delta x$$

and

$$(10d) \quad u_m^n = u(x_m, t_n) \quad .$$